

MODIFIED GHS COLLISION MODEL COMPARED WITH VHS MODEL

M. N. Macrossan and C. R. Lilley

School of Engineering
The University of Queensland
St Lucia, Brisbane, AUSTRALIA Q 4072
m.macrossan@uq.edu.au, c.lilley@uq.edu.au

The generalized hard sphere collision model (GHS) was introduced by Hassan and Hash [2]. It is generalization of the Sutherland collision model proposed by Kuscer [4]. At low temperatures, where the attractive intermolecular forces are important, the GHS collision model produces a more accurate variation of viscosity with temperature than the standard Variable Hard Sphere (VHS) collision model [1]. In spite of this, the GHS model remains virtually unused because of its great computational expense compared to the VHS model. A slight modification of the GHS model, described in [5], makes it no more than 5-15% more computationally expensive than the VHS model. We compare the GHS and VHS models for a blunt body flow with $T_0 = 1300$ K.




Fig. 1. Flat plate normal to flow. T contours and velocities. T_{wall} , front and back: 1300 K and 500 K.




Fig. 2. Modified GHS and VHS viscosity, compared with viscosity recommended for argon [3].

We calculate the supersonic flow of Argon, with a freestream temperature of 100 K, around a flat plate normal to the freestream (see Fig. 1). The wall temperatures were 1300 K (front) 500 K (rear), with diffuse reflection. The viscosity for the modified GHS and VHS models used here are compared in Fig. 2 with the recommended argon viscosity [3], over the range of temperatures in the flow. The temperature in the wake region is $\cong 500$ K, so the viscosity of the two models differs most there (see Fig. 2). This leads to small differences in the temperature field and the size of the re-circulation region. The GHS model required only 5% more CPU.

References

- [1] G. A. Bird, *Molecular Gas Dynamics and the Direct Simulation of Gas Flows*, Clarendon Press, 1994.
- [2] H. A. Hassan and D. B. Hash, "A Generalized Hard-Sphere Model for Monte-Carlo Simulation", *Physics Fluids A*, **5**, 738-744, 1993.
- [3] J. Kestin, K. Knierim, E. A. Mason, B. Najafi, S. T. Ro and Waldman, "Equilibrium Transport Properties of the Noble Gases and their Mixtures at Low Density", *J. Phys. Chem. Ref. Data*, **13**, 229-303, 1984
- [4] I. Kuscer, "A model for Rotational Energy Exchange in Polyatomic Gases", *Physica*, **158**, 784-764, 1989.
- [5] M. N. Macrossan and C. R. Lilley, "Modified Generalized Hard Sphere Collision Model for DSMC calculations", *Journal of Thermophysics Heat Trans*, **17**, 289-291, 2003.